



JAMDA

journal homepage: [www.jamda.com](http://www.jamda.com)

## Original Study

## Associations Between Daily Nurse Staffing Levels and Daily Hospitalizations and ED Visits in Nursing Homes

Dana B. Mukamel PhD<sup>a,\*</sup>, Debra Saliba MD, MPH<sup>b,c,d</sup>, Heather Ladd MS<sup>a</sup>, R. Tamara Konetzka PhD<sup>e</sup><sup>a</sup>Department of Medicine, Division of General Internal Medicine, iTEQC Research Program, University of California, Irvine, Irvine, CA, USA<sup>b</sup>University of California Los Angeles Borun Center at David Geffen School of Medicine, Los Angeles, CA, USA<sup>c</sup>Veterans Administration GRECC, Los Angeles, CA, USA<sup>d</sup>RAND Health, Santa Monica, CA, USA<sup>e</sup>Department of Public Health Sciences, The University of Chicago, Chicago, IL, USA

## ARTICLE INFO

## Article history:

Received 28 April 2022

Received in revised form

27 June 2022

Accepted 30 June 2022

## Keywords:

Nursing homes  
staffing  
hospitalizations  
ED visits  
quality

## ABSTRACT

**Objectives:** Although many prior studies have shown that high average levels of nurse staffing in nursing homes are associated with fewer hospitalizations, some studies have not, suggesting that the average nursing level may mask a more complex relationship. This study examines this issue by investigating the associations of daily staffing patterns and daily hospitalizations and emergency department (ED) visits. **Design:** Retrospective analyses of national Payroll Based Journal (PBJ) staffing data merged with the Minimum Data Set.

**Setting and Participants:** A total of 15,718 nursing homes nationally reporting PBJ data during 2017–2019, their staff, and residents.

**Methods:** We estimated facility-day-level models as conditional facility fixed-effect Poisson regressions with robust standard errors. The dependent variables were daily numbers of hospitalization and ED visits and the independent variables of interest were the number of registered nurse (RN), licensed practical nurse (LPN), and certified nurse assistant (CNA) hours on the same and prior days.

**Results:** The daily number of hospital transfers averaged 0.28 (SD 0.21). Daily total direct-care staffing hours averaged 288.7 (SD 188.2), with RNs accounting for 35.0, LPNs for 68.7, and CNAs for 185.0. Higher staffing was associated with more hospitalizations on the concurrent day. Higher staffing on the day prior was associated with fewer hospitalizations. The effect size was larger for RNs and LPNs (same day = ~2%; prior day = approximately –0.7% to –0.9%) than for CNAs (same day <1%; prior day < –0.5%). ED visits not leading to hospitalizations, and analyses for subsamples exhibited similar findings.

**Conclusions and Implications:** Our findings suggest that staff can address developing problems and prevent admissions the next day and identify emergent problems and hospitalize the same day. They also underscore the complex array of nursing home factors involved in hospitalization and ED visits, including the influence of daily staffing variation, suggesting the need for further research to better understand the associations between staffing and appropriate resident transfers to the hospital or the ED, and the potential implications for quality metrics in these domains.

© 2022 The Authors. Published by Elsevier Inc. on behalf of AMDA – The Society for Post-Acute and Long-Term Care Medicine. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Research reported in this publication was supported by the National Institute on Aging of the National Institutes of Health under Award Number R01AG066742. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

The authors declare no conflicts of interest.

\* Address correspondence to Dana B. Mukamel, PhD, iTEQC Research Program, Department of Medicine, Division of General Internal Medicine, University of California, Irvine, 100 Theory, Suite 120, Irvine, CA 92617–3056, USA.

E-mail address: [dmukamel@hs.uci.edu](mailto:dmukamel@hs.uci.edu) (D.B. Mukamel).

<https://doi.org/10.1016/j.jamda.2022.06.030>

1525–8610/© 2022 The Authors. Published by Elsevier Inc. on behalf of AMDA – The Society for Post-Acute and Long-Term Care Medicine. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Nursing homes are expected to provide most of the medical care their residents require in-house and minimize hospitalizations. Although some conditions, like acute myocardial infarction, almost always require hospitalization, many not only can be treated in the nursing home, but hospitalizing the resident may actually increase the risk of iatrogenic events and discontinuity of care contributing to further deterioration in the resident's condition, functional decline, and delirium.<sup>1–3</sup> Therefore, a low hospitalization rate has long been

considered a marker of high-quality nursing home care. Measures of appropriate hospitalizations were included in the Centers for Medicare and Medicaid Services (CMS) nursing home value-based payment demonstration<sup>4</sup> and measures of hospitalizations are included as quality indicators in the Nursing Home Care Compare report card published by CMS.<sup>5,6</sup>

Factors influencing hospitalizations have been studied extensively, with many studies examining staffing levels as explanatory variables, because staffing, whether registered nurses (RNs), licensed practical nurses (LPNs), or certified nurse assistants (CNAs), provide direct day-to-day patient care and are likely to be intimately involved in the care processes and decisions leading to hospitalizations. These studies, summarized by several recent reviews,<sup>7–12</sup> revealed mixed evidence about the relationship between staffing and hospitalization, with some failing to identify the expected association of higher staffing levels with lower hospitalization rates. The authors of these reviews identified a set of common limitations as likely reasons for the failure to find the expected associations between staffing and hospitalizations. These limitations include reliance on staffing data from the CMS Online Survey Certification and Reporting (OSCAR) System or the system that replaced it, the Certification and Survey Provider Enhanced Reporting (CASPER), both of which collect staffing data once a year and may be inaccurate,<sup>13,14</sup> the cross-sectional nature of most studies, and the lack of statistical control for other variables that might be important, such as presence of other providers paid by the facility (eg, nursing administrators, nurse practitioners, medical directors), and inability to distinguish between employed and contract staff.

We present analyses that address some of these prior limitations by using a new dataset, the Payroll Based Journal (PBJ), which provides daily staffing data based on each nursing home's payroll information, and is more accurate and more detailed than prior data. We take the novel approach of analyzing day-to-day patterns of hospitalizations as they relate to the daily staffing levels of RNs, LPNs, and CNAs for all nursing homes in the United States, as well as facilities at the top and bottom of the staffing distribution, offering a complementary perspective to prior studies. We test the hypothesis that the observed associations between average higher staffing and fewer hospitalizations will also be observed between same-day and prior-day staffing and hospitalizations.

## Methods

### Sample

The sample included all 15,718 Medicare- and Medicaid-certified nursing homes in the country during the period January 2017 through August 2019. The analysis for emergency department (ED) visits not followed by a hospitalization was based on outpatient claims files, which are limited to Medicare Fee-for-Service (FFS) residents and included 15,608 nursing homes (99.3%).

### Data Sources

The PBJ includes the number of hours paid daily for each staff type, including RNs, LPNs, CNAs, therapy staff, physicians, administrators, and others, by nursing home. It also provides daily patient census.

We merged the PBJ data with resident-level data, calculated from the Minimum Data Set (MDS) 3.0, at the facility-day level. The MDS includes assessments for all nursing home residents with information about age, gender, case mix [Resource Utilization Group (RUG) IV], cognitive impairment, Alzheimer or related dementias (ADRD) diagnoses, and dates of hospitalization and death. For some analyses we also used the Medicare outpatient claims for ED visits not followed by a hospital admission. We also merged in the Master Beneficiary Summary File, to identify residents enrolled in Medicare Advantage during their

nursing home stays, the Nursing Home Five-Star Quality Rating data, and facility characteristics reported in Long-Term-Care Focus.<sup>15</sup>

### Analyses

The study was approved by the University of California, Irvine, institutional review board.

Analyses were conducted at the facility-day level. Time-dependent variables were calculated for each nursing home, as counts or averages for each day.

### Dependent Variables

The dependent variables were the count of daily discharges from the nursing home to an acute care or psychiatric hospital as reported on the MDS discharge or transfer to the ED not followed by a hospital admission as reported on the Outpatient claim.

### Daily-Varying Independent Variables

There were 6 independent variables of interest: RN, LPN, and CNA hours on the transfer day to the hospital (or the ED) and RN, LPN, and CNA hours on the day before transfer. For example, when examining the number of transfers on July 1, we measure staffing on that same day (July 1) and on the preceding day (June 30).

Daily control variables included percent male residents in the facility on the day of transfer; percent residents who were younger than 65, 65 to 74, and 75 to 84, with 85+ as reference, and percentage of residents enrolled in Medicare Advantage on the transfer day. Case mix on the day of transfer was controlled by 66 variables. Each variable corresponds to 1 of the 66 RUGs categories and indicates the number of residents classified into that RUG on that date based on their most recent MDS assessment. Also included were a set of monthly indicator variables controlling for time trends.

### Stratification Variables for Subanalyses

We performed stratified analyses to examine hypotheses about different patterns of associations between staffing and hospitalizations in facilities with specific characteristics, as follows: (1) facility quality measured by the 5-Star Quality Measures (QMs) ratings and the 5-Star Survey ratings published in Nursing Home Care Compare and averaged over the study period; (2) average total direct-care nursing (sum of RNs, LPNs, and CNAs) hours over the study period; (3) average hospitalizations over the study period; (4) for-profit and nonprofit ownership; (5) percentage of residents covered by Medicare; and (6) percentage of residents with ADRD and cognitive impairment.

### Estimated Models

We estimated a facility-day-level model, with hospitalization on each day as the dependent variable and nursing hours by type (RN, LPN, and CNA) on the same day and on the day prior as the 6 independent variables of interest, controlling for all other variables described previously: age, sex, case-mix count, all of them calculated for the transfer day, and the month. The model was estimated as a conditional facility fixed-effect, Poisson regression, with robust standard errors. The facility fixed-effects approach controls for all time-invariant differences between nursing homes (eg, ownership).

We estimated a base case model on the full sample. We then stratified the sample by facility characteristics as described previously, and estimated models for subsamples defined by the bottom and top 30th percentiles of the distribution for each characteristic. We also

**Table 1**  
Descriptive Statistics

	Number of Nursing Homes	Mean	SD
Dependent Variables: Measured Daily			
Discharges from nursing home to acute care or psychiatric hospital	15,718	0.28	0.21
ED visits not followed by a hospitalization	15,608	0.11	0.07
Independent Variables: Nursing Homes and Resident Variables Measured Daily			
RN hours per day	15,718	35.0	33.3
Salaried	15,718	34.0	32.2
Contract	15,718	1.0	5.6
LPN hours per day	15,718	68.7	48.8
Salaried	15,718	66.4	47.0
Contract	15,718	2.3	7.6
CNA hours per day	15,718	185.0	124.8
Salaried	15,718	180.0	121.0
Contract	15,718	5.0	17.9
Total direct-care nurse staffing - per day	15,718	288.7	188.2
Director of nursing (RN) per day	15,718	5.0	1.7
RN Administrator per day	15,718	11.7	12.5
LPN administrator hours per day	15,718	6.0	8.7
Nurse aide in training hours per day	15,718	2.6	7.3
Medication aide hours per day	15,718	6.6	14.8
Medical director hours per day	15,718	0.4	0.7
Other MD hours per day	15,718	0.2	2.0
Nurse practitioner hours per day	15,718	0.3	1.2
Daily resident census	15,718	85.2	52.8
% of residents in Medicare Advantage	15,718	26.1	19.0
% of residents who are male	15,718	35.7	11.8
% of residents younger than 65	15,718	15.8	14.7
% of residents aged 65–74	15,718	18.5	8.0
% of residents aged 75–84	15,718	27.0	6.8
% of residents aged 85 and older	15,718	38.7	18.0
Facility Characteristics Measured Quarterly or Annually			
Average QMs 5-star rating (1–5)	13,422	3.8	1.0
Average survey 5-star rating (1–5)	13,455	2.8	1.1
% of residents with Medicare payer	14,960	13.3	13.4
% of residents with Medicaid payer	14,960	60.1	23.5
% of residents with ADRD and cognitive impairment diagnoses	15,718	47.7	16.2
	Number of Nursing Homes		Percent
For-profit ownership	10,479		70.1
Nonprofit ownership	4481		30.0

estimated a full sample model that separated employed and contract nursing staff and a full sample model that added other personnel types (if salaried by the facility) including hours of director of nursing-RN, RN administrators, LPN administrators, medical directors, other MDs, nurse practitioners, nurse aides in training, and medication aides.

We present the regression results for the staffing variables as a change in the percentage of daily hospitalizations associated with a 1 full-time equivalent (ie, 8 hours) increase in the staffing variable, holding all other variables constant. (To calculate this percentage, we used the formula  $\% \Delta(y|x) = 100 * (\exp(\beta * \Delta x))$ , where  $\% \Delta E(y|x)$  = expected % change in daily hospitalizations.) The *P* values reported are for the actual coefficients.

### Sensitivity Analyses

To test the robustness of our findings to different assumptions we performed several sensitivity analyses:

- To account for possible mismatches in dates between the hospital and ED data and the nursing home data, we estimated models in which transfer was matched more liberally, within 1 day of the discharge from the nursing home.
- To investigate sensitivity to the number of prior days' staffing included, we estimated models with (1) no variables for staffing on prior days, and (2) 2 variables for staffing 1 and 2 days prior.


- To test the sensitivity of our stratified analyses to the definition of the strata threshold, all analyses were repeated with thresholds set to 85% and 15%.
- To test whether our results are driven by the known phenomenon of lower staffing on weekends<sup>16</sup> or on Mondays and Fridays (as observed in our data, see later in this article), we performed an analysis limited to hospitalizations during midweek only (Tuesdays, Wednesdays, and Thursdays).

### Results

Table 1 presents descriptive statistics for the analytic sample. The average daily number of hospital transfers was 0.28 (SD 0.21) and transfers to the ED not followed by a hospitalization was 0.11 (SD 0.07). Average daily total direct-care staffing hours (including employed and contract staff) was 288.7 (SD 188.2). Of those, RNs accounted for 35.0, LPNs for 68.7, and CNAs for 185.0 hours, mostly accounted for by employed staff.

Figure 1 depicts daily staffing variations. It shows, separately for RNs, LPNs, and CNAs, the average number of hours per resident-day for each day of the week as a percent of the number of hours per resident-day averaged over the week, thus identifying days with staffing above and below average. All 3 staff types exhibit the same pattern, with the highest in midweek—Tuesday through Thursday—with approximately 2 to 5 percentage points decline on Friday and Monday, and the largest decline on the weekend, of more than 10 percentage points from the midweek high. The biggest decline is for

STAFFING	Avg RN Hours	Avg LPN Hours	Avg CNA Hours	As percent of weekly avg	Avg Daily Hosp Rate
	Per Res Day RNs	Per Res Day LPNs	Per Res Day CNAs		
Sunday	90.8%	92.3%	93.0%	Sunday	80.6%
Monday	102.1%	101.3%	100.7%	Monday	110.6%
Tuesday	104.7%	103.7%	103.2%	Tuesday	108.6%
Wednesday	105.1%	104.5%	104.0%	Wednesday	107.3%
Thursday	104.0%	103.6%	103.3%	Thursday	105.6%
Friday	101.5%	101.4%	100.9%	Friday	106.6%
Saturday	91.7%	93.2%	94.8%	Saturday	80.7%



\*Note: The weekly average is calculated separately for each of the 4 variables depicted in the chart.

The Y scale shows how the value of the variable on any given day deviates from the weekly average for that variable.

Example: On Sunday, hospitalizations are 81% of the hospitalization weekly average, RNs are at 91% of the RN weekly average, LPNs are at 95.5% of the LPN weekly average, and CNAs are at 96% of the CNA weekly average.

**Fig. 1.** Staffing per resident-day and hospitalizations by day of the week as percent of the weekly average. The weekly average is calculated separately for each of the 4 variables depicted in the chart. Example: On Sunday, hospitalizations are 81% of the hospitalization weekly average, RNs are at 91% of the RN weekly average, LPNs are at 95.5% of the LPN weekly average, and CNAs are at 96% of the CNA weekly average.

RNs and the smallest for CNAs. The figure also shows the daily variation in hospitalizations as percent of its weekly average. Hospitalizations follow a similar pattern to that of staffing, with a lower percent during the weekend compared with weekdays. The range of the differences between weekdays and weekend is larger for hospitalizations than for staffing, at approximately 25–30 percentage points.

Table 2 presents the results of the regression models, examining the relationships between staffing and hospitalizations for the base case, which includes the full sample, all nursing homes and all residents, followed by results of 2 models with different staffing specifications and then the stratified models. The table presents, for each model, the daily percent change in number of hospitalizations for a 1 full-time staff (FTE) increase. The full base case model is provided in Supplementary Table 1.

All models exhibit the same general findings. On the same day as the hospitalization, more staff is associated with more hospitalizations, typically with a stronger effect for RNs and LPNs, often approximately 2%, compared with CNAs, typically with less than 1% and often less than 0.5%. More staff on the day prior is associated with fewer hospitalizations, again mostly with stronger effects for RNs and LPNs, mostly between  $-0.7\%$  and  $-0.9\%$  and smaller effects for CNAs, less than  $-0.5\%$ .

All effects are significant at the 0.001 level, except for 2 models. The model with separate covariates for employed and contract staff

has very low previous day effects with  $P$  values of 0.02, 0.7, and 0.4 for contract RNs, LPNs, and CNAs, respectively. The model with additional types of non-nursing staff has a very low same day and nonsignificant ( $P = .15$ ) effect for CNAs.

The sensitivity analyses described previously have similar findings. A model without any lags shows a positive association between staffing and hospitalization on the same day. Adding 2-days lags show a positive same-day effect and negative effects 1 and 2 days before the transfer. Supplementary Table 2 shows summary results for dyads defined with different thresholds and for the midweek days sample.

## Discussion

In this study, we evaluated the association between daily staffing and hospitalization and found that it depends on the day of transfer to the hospital. High staffing was protective against hospitalization on the following day, but high staffing measured on the concurrent day with the hospitalization was associated with more hospitalizations. These findings were very robust, observed not only in the full sample and base case, but also in all other model specifications, when estimated separately for employed and contract staff, when non-nursing staff were added to the model, and when the model was estimated on stratified samples.

**Table 2**  
**Summary of Findings: Base Case and Stratified Analyses—Percent Change in Daily Hospitalizations Due to an Increase of a 1 Full-Time Equivalent Staff Position\***

	Same Day			Previous Day		
	RNs	LPNs	CNAs	RNs	LPNs	CNAs
Base case, full sample	<b>2.3% [2.1, 2.4] (&lt;.001)</b>	<b>2.1% [2.0, 2.2] (&lt;.001)</b>	<b>0.6% [0.5, 0.7] (&lt;.001)</b>	<b>-0.8% [-0.9, -0.7] (&lt;.001)</b>	<b>-0.8% [-0.9, -0.7] (&lt;.001)</b>	<b>-0.4% [-0.4, -0.3] (&lt;.001)</b>
Models with different specifications, full sample:						
(1) Separate staffing variables for employed and contract staff by type in same model, full sample						
Employed staff	<b>2.3% [2.2, 2.4] (&lt;.001)</b>	<b>2.1% [2.0, 2.3] (&lt;.001)</b>	<b>0.6% [0.6, 0.7] (&lt;.001)</b>	<b>-0.8% [-0.9, -0.7] (&lt;.001)</b>	<b>-0.8% [-0.9, -0.8] (&lt;.001)</b>	<b>-0.4% [-0.4, -0.3] (&lt;.001)</b>
Contract staff	<b>1.7% [1.3, 2.1] (&lt;.001)</b>	<b>1.4% [1.2, 1.6] (&lt;.001)</b>	<b>0.3% [0.2, 0.5] (&lt;.001)</b>	-0.4% [-0.7, -0.1] (.017)	-0.0% [-0.2, 0.1] (.698)	-0.1% [-0.2, 0.1] (.383)
(2) Adding variables for non-nursing FTE providers to the model, <sup>†</sup> full sample	<b>1.5% [1.3, 1.6] (&lt;.001)</b>	<b>1.5% [1.4, 1.6] (&lt;.001)</b>	0.0% [-0.0, 0.1] (.149)	<b>-0.3% [-0.4, -0.2] (&lt;.001)</b>	<b>-0.2% [-0.3, -0.1] (&lt;.001)</b>	<b>-0.1% [-0.2, -0.1] (&lt;.001)</b>
(3) NH Quality measured by 5-Star QMs stratification						
High average quality 5-star rating (≥4.75)	<b>2.4% [2.1, 2.7] (&lt;.001)</b>	<b>2.1% [1.8, 2.5] (&lt;.001)</b>	<b>0.5% [0.3, 0.7] (&lt;.001)</b>	<b>-0.9% [-1.2, -0.6] (&lt;.001)</b>	<b>-0.8% [-1.0, -0.6] (&lt;.001)</b>	<b>-0.3% [-0.4, -0.1] (.001)</b>
Low average quality 5-star rating (<2.5)	<b>2.0% [1.6, 2.4] (&lt;.001)</b>	<b>2.0% [1.7, 2.4] (&lt;.001)</b>	<b>0.8% [0.7, 1.0] (&lt;.001)</b>	<b>-0.6% [-0.9, -0.3] (&lt;.001)</b>	<b>-0.8% [-1.0, -0.6] (&lt;.001)</b>	<b>-0.3% [-0.5, -0.1] (.001)</b>
(4) NH quality measured by 5-star survey stratification						
High average survey 5-star rating (≥4)	<b>2.4% [2.0, 2.7] (&lt;.001)</b>	<b>2.0% [1.6, 2.3] (&lt;.001)</b>	<b>0.5% [0.2, 0.7] (0.003)</b>	<b>-0.7% [-0.9, -0.4] (&lt;.001)</b>	<b>-0.7% [-1.0, -0.5] (&lt;.001)</b>	<b>-0.3% [-0.5, -0.1] (.001)</b>
Low average survey 5-star rating (<2)	<b>2.3% [2.1, 2.5] (&lt;.001)</b>	<b>2.2% [2.0, 2.4] (&lt;.001)</b>	<b>0.7% [0.6, 0.8] (&lt;.001)</b>	<b>-0.9% [-1.1, -0.8] (&lt;.001)</b>	<b>-0.9% [-1.0, -0.8] (&lt;.001)</b>	<b>-0.4% [-0.5, -0.3] (&lt;.001)</b>
(5) Total nursing level stratification						
High average nursing level (above 70 <sup>th</sup> percentile)	<b>2.3% [2.1, 2.5] (&lt;.001)</b>	<b>2.0% [1.8, 2.2] (&lt;.001)</b>	<b>0.4% [0.2, 0.5] (&lt;.001)</b>	<b>-0.9% [-1.1, -0.7] (&lt;.001)</b>	<b>-0.8% [-1.0, -0.7] (&lt;.001)</b>	<b>-0.2% [-0.3, -0.1] (&lt;.001)</b>
Low average nursing level (30 <sup>th</sup> percentile and below)	<b>2.1% [1.9, 2.4] (&lt;.001)</b>	<b>2.2% [2.0, 2.5] (&lt;.001)</b>	<b>0.9% [0.8, 1.1] (&lt;.001)</b>	<b>-0.7% [-0.9, -0.5] (&lt;.001)</b>	<b>-0.9% [-1.0, -0.7] (&lt;.001)</b>	<b>-0.5% [-0.6, -0.4] (&lt;.001)</b>
(6) Hospitalization-level stratification						
High hospitalization rate (above 70 <sup>th</sup> percentile)	<b>2.4% [2.2, 2.6] (&lt;.001)</b>	<b>2.3% [2.1, 2.5] (&lt;.001)</b>	<b>0.7% [0.5, 0.8] (&lt;.001)</b>	<b>-0.9% [-1.1, -0.8] (&lt;.001)</b>	<b>-0.9% [-1.1, -0.8] (&lt;.001)</b>	<b>-0.5% [-0.6, -0.3] (&lt;.001)</b>
Low hospitalization rate (30 <sup>th</sup> percentile and below)	<b>2.0% [1.7, 2.4] (&lt;.001)</b>	<b>1.6% [1.3, 1.9] (&lt;.001)</b>	<b>0.4% [0.3, 0.5] (&lt;.001)</b>	<b>-0.6% [-0.7, -0.4] (&lt;.001)</b>	<b>-0.5% [-0.7, -0.3] (&lt;.001)</b>	<b>-0.2% [-0.3, -0.1] (&lt;.001)</b>
(7) Nonprofit vs for-profit nursing homes						
Nonprofit nursing homes	<b>1.9% [1.7, 2.1] (&lt;.001)</b>	<b>1.4% [1.2, 1.6] (&lt;.001)</b>	<b>0.5% [0.4, 0.6] (&lt;.001)</b>	<b>-0.7% [-0.9, -0.5] (&lt;.001)</b>	<b>-0.6% [-0.8, -0.5] (&lt;.001)</b>	<b>-0.2% [-0.3, -0.1] (&lt;.001)</b>
For-profit nursing homes	<b>2.5% [2.3, 2.6] (&lt;.001)</b>	<b>2.4% [2.3, 2.5] (&lt;.001)</b>	<b>0.7% [0.6, 0.8] (&lt;.001)</b>	<b>-0.8% [-1.0, -0.7] (&lt;.001)</b>	<b>-0.8% [-0.9, -0.8] (&lt;.001)</b>	<b>-0.5% [-0.5, -0.4] (&lt;.001)</b>
(8) High vs low Medicare census nursing homes						
High Medicare nursing homes (above 70 <sup>th</sup> percentile)	<b>2.4% [2.2, 2.6] (&lt;.001)</b>	<b>2.2% [2.0, 2.4] (&lt;.001)</b>	<b>0.7% [0.5, 0.8] (&lt;.001)</b>	<b>-0.8% [-1.0, -0.7] (&lt;.001)</b>	<b>-0.9% [-1.1, -0.8] (&lt;.001)</b>	<b>-0.4% [-0.6, -0.3] (&lt;.001)</b>
Low Medicare nursing homes (30 <sup>th</sup> percentile and below)	<b>2.0% [1.8, 2.3] (&lt;.001)</b>	<b>1.8% [1.6, 2.0] (&lt;.001)</b>	<b>0.6% [0.5, 0.7] (&lt;.001)</b>	<b>-0.6% [-0.8, -0.4] (&lt;.001)</b>	<b>-0.6% [-0.8, -0.5] (&lt;.001)</b>	<b>-0.4% [-0.5, -0.3] (&lt;.001)</b>
(9) Nursing homes with high and low percentage of residents with ADRD and cognitive impairment						
High ADRD (above 70 <sup>th</sup> percentile)	<b>2.1% [1.9, 2.3] (&lt;.001)</b>	<b>1.9% [1.7, 2.2] (&lt;.001)</b>	<b>0.5% [0.4, 0.6] (&lt;.001)</b>	<b>-0.7% [-0.8, -0.6] (&lt;.001)</b>	<b>-0.7% [-0.9, -0.6] (&lt;.001)</b>	<b>-0.3% [-0.4, -0.2] (&lt;.001)</b>
Low ADRD (30 <sup>th</sup> percentile and below)	<b>2.7% [2.5, 2.8] (&lt;.001)</b>	<b>2.4% [2.3, 2.6] (&lt;.001)</b>	<b>0.8% [0.7, 0.9] (&lt;.001)</b>	<b>-1.0% [-1.1, -0.8] (&lt;.001)</b>	<b>-1.0% [-1.1, -0.9] (&lt;.001)</b>	<b>-0.6% [-0.7, -0.5] (&lt;.001)</b>
(10) Residents transferred to the ED but not followed by a hospitalization						
Based on outpatient claims files that include FFS residents only	<b>1.4% [1.3, 1.6] (&lt;.001)</b>	<b>1.5% [1.3, 1.6] (&lt;.001)</b>	<b>0.1% [0.1, 0.2] (&lt;.001)</b>	<b>-0.4% [-0.5, -0.3] (&lt;.001)</b>	<b>-0.4% [-0.5, -0.3] (&lt;.001)</b>	<b>-0.1% [-0.1, -0.0] (.023)</b>

Bold values are statistically significant.

\*Each line reports findings from a separate regression model that also controls for age, gender, case mix, seasonality, and facility fixed effects. Models differ by either staffing variables (eg, salaried vs contract), or facility characteristics (eg, low quality vs high quality).

<sup>†</sup>Additional providers included director of nursing-RN, RN administrator, LPN administrator, medical director, other MD, nurse practitioner, nurses aid in training, medication aid.

Although the protective effect of high staffing on hospitalization has been observed in many previous studies,<sup>7–12</sup> evidence for high staffing increasing the risk of hospitalization for nursing home residents is novel, to our knowledge. We believe that these unexpected findings are because for the first time we are able to analyze daily staffing data. All prior studies of the associations between staffing and hospitalizations relied on staffing data averaged over large periods of time, which likely masked this phenomenon.

What might be the explanation for these 2 different associations between staffing and hospitalizations? The same-day increased hospitalization with higher staffing is likely due to staff's increased ability to monitor residents and identify those residents who may require immediate transfer. We note that the same-day effect is more than 3 times larger for RNs and LPNs than for CNAs. As the former are the ones likely to perform assessments and make transfer recommendations, this offers further support for the explanation. The protective effect of the prior-day high staffing on hospitalization may also be explained by staff having increased ability to monitor and identify residents with clinical issues requiring interventions. In these cases, the interventions available in-house might avert further decline and prevent the need for hospitalization when given time. Hence, increased staffing coupled with more time (ie, the additional day or two) mean that some hospitalizations can be prevented, such that on net, the effect of increased staffing is to lower hospitalizations.

Comparing the magnitude of the associations across strata dyads in many cases offers further support for these explanations. For example, we find higher effect sizes for employed staff who know the residents better and are, therefore, more likely to ascertain their needs than contract staff. We find higher effect sizes for RNs in nursing homes with higher 5-star ratings for both the QMs and the Survey, as well as facilities that have higher total nursing levels. The high Medicare facilities, which have more post-acute patients who are more likely to have higher patient acuity and skilled needs that might require rehospitalizations also exhibit larger effects than nursing homes with fewer Medicare patients. Nursing homes with higher census of patients with ADRD diagnoses have lower effect size compared with nursing homes with fewer patients with these diagnoses, possibly reflecting a more judicious approach toward hospital transfer for patients with ADRD, as high-quality care practices would suggest. Although another less benign possibility is that residents with ADRD are less likely to express themselves and staff may not recognize the need to hospitalize as often.

We note that the short-term patterns we observe are not likely to be due to long-term staff shortages. Our sensitivity analysis comparing nursing homes with high and low average staffing found the same staffing-to-hospitalization associations among both, suggesting that long-term staffing differences do not explain away these patterns. We also note that we did not control for turnover because it is likely one of the mechanisms for daily staffing variations, and including it would mask the effect of interest. Future studies examining factors leading to daily staffing patterns should include turnover.

Several limitations of this study should be mentioned. Because of data limitation, our estimates of ED transfers were limited to FFS residents only. Second, the associations we found are not necessarily causal; however, we used a strong facility fixed-effects design to control for confounding by all time-invariant facility-level characteristics. Third, our staffing measures are at the facility level and we cannot identify the staff hours received by individual residents or groups of residents, such as those with ADRD, or whether or not residents had private duty nurses or family involvement. Finally, these quantitative data cannot establish the mechanisms driving the associations we find. For example, we did not have data about nurses' training to start intravenous lines in the nursing home, manage total parenteral nutrition or tube feeding. Furthermore, we focused on staffing only and had no information about availability of other

resources, such as laboratory and x-ray results within 4 to 8 hours, specialists, hospice services, and 7 days-per-week therapy services, all of which are considered important for preventing hospitalizations.<sup>3</sup>

## Conclusions and Implications

This is the first study, to our knowledge, to examine short-term patterns of staffing in nursing homes and their associations with hospitalizations and ED transfers not followed by a hospitalization. The richness of the PBJ data allowed us to examine daily staffing patterns vis-à-vis daily outcomes, and revealed patterns that have not been observed before. Our findings reaffirm the importance of staffing to hospitalization and ED outcomes, even when inspected on a microlevel, and offer a new perspective for understanding the roles of different staffing types. Specifically, our findings suggest that staffing patterns, especially for RNs and LPNs, play a key role in assessing and triaging residents, both in identifying those residents who require immediate hospitalization, thus contributing to the higher number of hospitalizations on the same day, and in identifying those in which hospitalizations can be averted if appropriate action is taken early enough.

An indirect implication of our study is that a simple count or rate of hospitalizations may not be a good marker for quality, and a more nuanced approach should be considered. The use of hospitalization and rehospitalization rates has become ubiquitous as a quality metric. At the same time, it has long been recognized that some hospitalizations are necessary, and that failing to transfer may be a sign of poor quality. Our study reinforces the need to remember this caveat and continue to refine quality metrics to try to disentangle appropriate and inappropriate transfers.

This study offers new observations about the associations between staffing and hospitalizations. Its strength is its reliance on national data. This, however, is also its weakness. It cannot get into the details of the underlying processes of care, which lead to the outcomes and associations we observe, and some of the questions we raised previously. Future studies, relying on other data sources and different methods, should address these questions, as well as the mechanisms leading to daily staffing variations.

## Acknowledgments

Debra Saliba is an employee of the Veterans Health Administration. The views presented here do not represent those of the Department of Veterans Affairs. The sponsor had no role except for funding.

## References

1. Creditor MC. Hazards of hospitalization of the elderly. *Ann Intern Med.* 1993; 118:219–223.
2. Dwyer R, Gabbe B, Stoelwinder JU, Lowthian J. A systematic review of outcomes following emergency transfer to hospital for residents of aged care facilities. *Age Ageing.* 2014;43:759–766.
3. Ouslander JG, Lamb G, Perloe M, et al. Potentially avoidable hospitalizations of nursing home residents: frequency, causes, and costs: [see editorial comments by Drs. Jean F. Wyman and William R. Hazzard, pp 760–761]. *J Am Geriatr Soc.* 2010;58:627–635.
4. CMS.gov. Nursing home value-based purchasing demonstration. 2009. Accessed November 24, 2021. <https://innovation.cms.gov/innovation-models/nursing-home-value-based-purchasing>
5. CMS.gov. Nursing homes - quality of resident care. Accessed November 24, 2021. <https://data.cms.gov/provider-data/topics/nursing-homes/quality-of-resident-care/#quality-of-resident-care-data-collection-periods>
6. Saliba D, Weimer DL, Shi Y, Mukamel DB. Examination of the new short-stay nursing home quality measures: rehospitalizations, emergency department visits, and successful returns to the community. *Inquiry.* 2018;55:46958018786816.
7. Laging B, Ford R, Bauer M, Nay R. A meta-synthesis of factors influencing nursing home staff decisions to transfer residents to hospital. *J Adv Nurs.* 2015; 71:2224–2236.

8. Backhaus R, Verbeek H, van Rossum E, Capezuti E, Hamers JP. Nurse staffing impact on quality of care in nursing homes: a systematic review of longitudinal studies. *J Am Med Dir Assoc*. 2014;15:383–393.
9. Spilsbury K, Hewitt C, Stirk L, Bowman C. The relationship between nurse staffing and quality of care in nursing homes: a systematic review. *Int J Nurs Stud*. 2011;48:732–750.
10. Grabowski DC, Stewart KA, Broderick SM, Coots LA. Predictors of nursing home hospitalization: a review of the literature. *Med Care Res Rev*. 2008;65:3–39.
11. Castle NG. Nursing home caregiver staffing levels and quality of care: a literature review. *J Appl Gerontol*. 2008;27:375–405.
12. Bostick JE, Rantz MJ, Flesner MK, Riggs CJ. Systematic review of studies of staffing and quality in nursing homes. *J Am Med Dir Assoc*. 2006;7:366–376.
13. Kash BA, Hawes C, Phillips CD. Comparing Staffing Levels in the Online Survey Certification and Reporting (OSCAR) system with the Medicaid cost report data: are differences systematic? *Gerontologist*. 2007;47:480–489.
14. Feng Z, Katz PR, Intrator O, Karuza J, Mor V. Physician and nurse staffing in nursing homes: the role and limitations of the Online Survey Certification and Reporting (OSCAR) system. *J Am Med Dir Assoc*. 2005;6:27–33.
15. LTC Focus, Brown University. Facts on Care in the US - Create Custom Reports on Long-Term Care. Accessed May 11, 2022. <https://ltafocus.org/>
16. Centers for Medicare and Medicaid Services. Design for care compare nursing home five-star quality rating system: technical users' guide-January 2022. Accessed March 31, 2022. <https://www.cms.gov/medicare/provider-enrollment-and-certification/certificationandcompliance/downloads/usersguide.pdf>

**Supplementary Table 1**

Full Regression Model of Base Case: Full Sample, Dependent Variable: Number of Hospitalizations from the Nursing Home

	Coefficient	Robust SE	z	P > z	95% CI		
Conditional fixed-effects Poisson regression							
Number of observations = 14,620,886							
Number of groups = 15,718							
Observations per group:							
min = 27							
avg = 930.2							
max = 1002							
Wald $\chi^2$ (110) = 11,866.77							
Log pseudolikelihood = -8,927,513.8							
Prob > $\chi^2$ = 0.0000							
(SE adjusted for clustering on id)							
	Coefficient	Robust SE	z	P > z	95% CI		
RN hours	0.002795	7.79E-05	35.87	<.001	0.002642	0.002947	
LPN hours	0.002603	6.94E-05	37.48	<.001	.002466	0.002739	
CNA hours	0.000774	4.62E-05	16.76	<.001	.000683	0.000864	
1 day lag RN hours	-0.00102	6.05E-05	-16.81	<.001	-.00114	-0.0009	
1 day lag LPN hours	-0.00101	5.14E-05	-19.69	<.001	-.00111	-0.00091	
1 day lag CNA hours	-0.00046	3.82E-05	-11.95	<.001	-.00053	-0.00038	
Percent male	0.003463	0.000207	16.73	<.001	.003058	0.003869	
Percent age 65 or less	0.005515	0.000368	14.98	<.001	.004794	0.006236	
pctage65_74	0.006117	0.000293	20.86	<.001	.005543	0.006692	
pctage75_84	0.004348	0.000237	18.38	<.001	.003884	0.004811	
Percent Medicare Advantage	-0.00173	0.000231	-7.5	<.001	-.00218	-0.00128	
Jan-17	-0.00582	0.017593	-0.33	.741	-.0403	0.028666	
Feb-17	0.154582	0.005393	28.67	<.001	.144012	0.165151	
Mar-17	0.174678	0.007983	21.88	<.001	.159032	0.190324	
Apr-17	0.128258	0.00882	14.54	<.001	.110971	0.145545	
May-17	0.115096	0.008917	12.91	<.001	.097618	0.132573	
Jun-17	0.099164	0.008807	11.26	<.001	.081904	0.116424	
Jul-17	0.065309	0.008853	7.38	<.001	.047958	0.082661	
Aug-17	0.087147	0.008749	9.96	<.001	.069999	0.104295	
Sep-17	0.090546	0.008594	10.54	<.001	.073701	0.107391	
Oct-17	0.096721	0.008598	11.25	<.001	.079868	0.113573	
Nov-17	0.099547	0.008659	11.5	<.001	.082575	0.116518	
Dec-17	0.152027	0.008327	18.26	<.001	.135707	0.168347	
Jan-18	0.231686	0.007924	29.24	<.001	.216155	0.247217	
Feb-18	0.174271	0.007959	21.9	<.001	.158672	0.18987	
Mar-18	0.142361	0.008027	17.74	<.001	.126629	0.158093	
Apr-18	0.122641	0.008195	14.97	<.001	.106579	0.138703	
May-18	0.093686	0.008257	11.35	<.001	.077503	0.109869	
Jun-18	0.078696	0.008139	9.67	<.001	.062744	0.094648	
Jul-18	0.070025	0.008036	8.71	<.001	.054275	0.085775	
Aug-18	0.083819	0.007951	10.54	<.001	.068236	0.099403	
Sep-18	0.078109	0.007946	9.83	<.001	.062536	0.093683	
Oct-18	0.091745	0.007728	11.87	<.001	.076599	0.10689	
Nov-18	0.07834	0.007808	10.03	<.001	.063037	0.093644	
Dec-18	0.094132	0.007611	12.37	<.001	.079215	0.109049	
Jan-19	0.141538	0.007398	19.13	<.001	.127038	0.156038	
Feb-19	0.142629	0.007009	20.35	<.001	.128893	0.156366	
Mar-19	0.138552	0.006835	20.27	<.001	.125157	0.151948	
Apr-19	0.113781	0.006824	16.67	<.001	.100407	0.127155	
May-19	0.085539	0.006785	12.61	<.001	.072242	0.098836	
Jun-19	0.061354	0.006765	9.07	<.001	.048094	0.074614	
Jul-19	0.058282	0.006559	8.89	<.001	.045427	0.071138	
Aug-19	0.038236	0.005584	6.85	<.001	.027291	0.04918	
Major RUG Group	RUG Code	Coefficient	Robust SE	z	P > z	95% CI	
Behavioral Symptoms and Cognitive Performance	ba1	0.006547	0.000531	12.33	<.001	0.005506	0.007588
	ba2	0.004166	0.001641	2.54	.011	0.00095	0.007381
	bb1	0.008505	0.000531	16	<.001	0.007464	0.009547
	bb2	0.003671	0.001543	2.38	.017	0.000646	0.006695
	ca1	0.010229	0.00088	11.62	<.001	0.008504	0.011954
Clinically Complex	ca2	0.011779	0.001703	6.92	<.001	0.008441	0.015117
	cb1	0.011881	0.000864	13.75	<.001	0.010187	0.013575
	cb2	0.016374	0.003438	4.76	<.001	0.009635	0.023113
	cc1	0.010164	0.000562	18.08	<.001	0.009062	0.011266
	cc2	0.010332	0.002008	5.15	<.001	0.006396	0.014267
	cd1	0.010293	0.000593	17.37	<.001	0.009131	0.011455
	cd2	0.010185	0.001986	5.13	<.001	0.006293	0.014077
	ce1	0.008007	0.000937	8.55	<.001	0.006171	0.009844
Extensive Services	ce2	0.008822	0.003826	2.31	.021	0.001324	0.01632
	es1	0.011822	0.001398	8.45	<.001	0.009081	0.014562
	es2	0.010938	0.001597	6.85	<.001	0.007807	0.014068
	es3	0.008712	0.001996	4.37	<.001	0.004801	0.012624

(continued on next page)



Supplementary Table 1 (continued)

Major RUG Group	RUG Code	Coefficient	Robust SE	z	P > z	95% CI		
Special Care High	hb1	0.010922	0.000974	11.22	<.001	0.009014	0.01283	
	hb2	0.008333	0.001727	4.82	<.001	0.004948	0.011718	
	hc1	0.009431	0.000878	1.74	<.001	0.00771	0.011152	
	hc2	0.006885	0.001626	4.23	<.001	0.003698	0.010072	
	hd1	0.009922	0.000776	12.79	<.001	0.008401	0.011443	
	hd2	0.006278	0.001619	3.88	<.001	0.003105	0.00945	
	he1	0.010329	0.001079	9.58	<.001	0.008214	0.012443	
	he2	0.006903	0.002076	3.33	.001	0.002835	0.010971	
Special Care Low	lb1	0.015033	0.001058	14.21	<.001	0.01296	0.017106	
	lb2	0.014114	0.003702	3.81	<.001	0.006859	0.021369	
	lc1	0.011452	0.000577	19.87	<.001	0.010322	0.012582	
	lc2	0.011764	0.002169	5.42	<.001	0.007513	0.016014	
	ld1	0.010339	0.000576	17.94	<.001	0.00921	0.011468	
	ld2	0.011236	0.001692	6.64	<.001	0.007919	0.014553	
	le1	0.009884	0.000764	12.93	<.001	0.008386	0.011382	
	le2	0.007856	0.002206	3.56	<.001	0.003532	0.01218	
Reduced Physical Functioning	pa1	0.00777	0.00052	14.95	<.001	0.006751	0.008788	
	pa2	0.008026	0.001431	5.61	<.001	0.005222	0.010831	
	pb1	0.007634	0.000539	14.15	<.001	0.006577	0.008691	
	pb2	0.008797	0.001105	7.96	<.001	0.006631	0.010963	
	pc1	0.008381	0.00045	18.64	<.001	0.0075	0.009263	
	pc2	0.00676	0.000747	9.05	<.001	0.005297	0.008224	
	pd1	0.0084	0.000413	2.32	<.001	0.007589	0.00921	
	pd2	0.00553	0.000874	6.33	<.001	0.003817	0.007243	
	pe1	0.006693	0.000671	9.98	<.001	0.005378	0.008008	
	pe2	0.004636	0.001325	3.5	<.001	0.002039	0.007233	
	Rehab and Rehab Plus Extensive	rha	0.008448	0.000604	13.98	<.001	0.007264	0.009633
		rhb	0.011513	0.000632	18.22	<.001	0.010274	0.012751
rhc		0.011504	0.000568	2.26	<.001	0.010391	0.012617	
rhl		0.016735	0.003125	5.35	<.001	0.010609	0.022861	
rhx		0.017419	0.002393	7.28	<.001	0.012729	0.022108	
rla		0.007349	0.000721	1.19	<.001	0.005935	0.008762	
rlb		0.008029	0.000986	8.14	<.001	0.006097	0.009961	
rlx		0.015218	0.004622	3.29	.001	0.006159	0.024276	
rma		0.007689	0.000466	16.49	<.001	0.006775	0.008603	
rmb		0.009195	0.00043	21.38	<.001	0.008352	0.010038	
rmc		0.009279	0.000433	21.43	<.001	0.008431	0.010128	
rml		0.013684	0.00393	3.48	<.001	0.005982	0.021386	
rmx		0.007214	0.003217	2.24	.025	0.000909	0.013519	
rua		0.010369	0.000443	23.42	<.001	0.009501	0.011236	
rub		0.009272	0.000424	21.86	<.001	0.008441	0.010103	
ruc		0.011112	0.000476	23.35	<.001	0.01018	0.012045	
rul		0.015571	0.002003	7.78	<.001	0.011646	0.019496	
rux		0.012584	0.002795	4.5	<.001	0.007106	0.018063	
rva	0.01164	0.00059	19.73	<.001	0.010484	0.012796		
rvb	0.012082	0.000608	19.86	<.001	0.01089	0.013274		
rvc	0.013594	0.000609	22.32	<.001	0.012401	0.014788		
rvl	0.025555	0.002493	1.25	<.001	0.020668	0.030441		
rvx	0.021725	0.001883	11.54	<.001	0.018034	0.025416		
Unknown		0.032939	0.002768	11.9	0	<0.001	0.038364	

**Supplementary Table 2**

Sensitivity Analysis Summary of Findings: Percent Change in Daily Hospitalizations Due to an Increase of a 1 Full-Time Equivalent Staff Position\*

	Same Day			Previous Day		
	RNs	LPNs	CNAs	RNs	LPNs	CNAs
Models Estimated on Stratified Samples						
Midweek days only						
NH Quality measured by 5 Star <u>QMs</u>	<b>1.0% [0.9, 1.1] (&lt;.001)</b>	<b>1.0% [0.9, 1.1] (&lt;.001)</b>	<b>0.0% [-0.0, 0.1] (0.698)</b>	<b>-0.5% [-0.6, -0.4] (&lt;.001)</b>	<b>-0.4% [-0.5, -0.3] (&lt;.001)</b>	-0.0% [-0.1, 0.0] (0.148)
High Average Quality	<b>2.5% [2.2, 2.7] (&lt;.001)</b>	<b>2.2% [1.9, 2.5] (&lt;.001)</b>	<b>0.5% [0.3, 0.7] (&lt;.001)</b>	<b>-0.9% [-1.1, -0.7] (&lt;.001)</b>	<b>-0.8% [-1.0, -0.6] (&lt;.001)</b>	<b>-0.3% [-0.4, -0.1] (&lt;.001)</b>
5-Star Rating (70 <sup>th</sup> percentile)						
Low Average Quality	<b>2.1% [1.9, 2.3] (&lt;.001)</b>	<b>2.1% [1.9, 2.3] (&lt;.001)</b>	<b>0.9% [0.8, 1.0] (&lt;.001)</b>	<b>-0.8% [-1.0, -0.6] (&lt;.001)</b>	<b>-0.9% [-1.1, -0.8] (&lt;.001)</b>	<b>-0.4% [-0.5, -0.3] (&lt;.001)</b>
5-Star Rating (30 <sup>th</sup> percentile)						
NH Quality measured by 5-Star <u>Survey</u>						
High Average Survey	<b>2.4% [2.1, 2.8] (&lt;.001)</b>	<b>2.2% [1.9, 2.5] (&lt;.001)</b>	<b>0.5% [0.3, 0.8] (&lt;.001)</b>	<b>-0.8% [-1.0, -0.6] (&lt;.001)</b>	<b>-0.8% [-1.1, -0.6] (&lt;.001)</b>	<b>-0.3% [-0.5, -0.2] (&lt;.001)</b>
5-Star Rating (70 <sup>th</sup> percentile)						
Low Average Survey	<b>2.3% [2.1, 2.6] (&lt;.001)</b>	<b>2.2% [2.0, 2.4] (&lt;.001)</b>	<b>0.7% [0.6, 0.8] (&lt;.001)</b>	<b>-0.9% [-1.1, -0.8] (&lt;.001)</b>	<b>-0.9% [-1.0, -0.8] (&lt;.001)</b>	<b>-0.4% [-0.5, -0.3] (&lt;.001)</b>
5-Star Rating (30 <sup>th</sup> percentile)						
Total nursing level variations						
High Average Nursing Level (85 <sup>th</sup> percentile)	<b>2.2% [1.9, 2.5] (&lt;.001)</b>	<b>1.8% [1.6, 2.1] (&lt;.001)</b>	<b>0.3% [0.1, 0.5] (&lt;.001)</b>	<b>-0.8% [-0.9, -0.6] (&lt;.001)</b>	<b>-0.7% [-0.9, -0.5] (&lt;.001)</b>	<b>-0.2% [-0.3, -0.1] (0.004)</b>
Low Average Nursing Level (15 <sup>th</sup> percentile)	<b>2.1% [1.7, 2.4] (&lt;.001)</b>	<b>2.3% [1.9, 2.7] (&lt;.001)</b>	<b>1.0% [0.7, 1.2] (&lt;.001)</b>	<b>-0.9% [-1.3, -0.6] (&lt;.001)</b>	<b>-0.9% [-1.2, -0.6] (&lt;.001)</b>	<b>-0.7% [-0.9, -0.5] (&lt;.001)</b>
Hospitalization-level variations						
High Hospitalization Rate (85 <sup>th</sup> percentile)	<b>2.5% [2.2, 2.7] (&lt;.001)</b>	<b>2.4% [2.2, 2.6] (&lt;.001)</b>	<b>0.7% [0.5, 0.8] (&lt;.001)</b>	<b>-1.0% [-1.2, -0.8] (&lt;.001)</b>	<b>-1.0% [-1.2, -0.9] (&lt;.001)</b>	<b>-0.5% [-0.6, -0.4] (&lt;.001)</b>
Low Hospitalization Rate (15 <sup>th</sup> percentile)	<b>1.9% [1.3, 2.5] (&lt;.001)</b>	<b>1.6% [1.2, 2.0] (&lt;.001)</b>	<b>0.5% [0.3, 0.6] (&lt;.001)</b>	<b>-0.3% [-0.6, -0.1] (0.013)</b>	<b>-0.2% [-0.4, 0.1] (0.226)</b>	<b>-0.3% [-0.5, -0.1] (0.001)</b>
High vs low Medicare census nursing homes						
High Medicare facilities (85 <sup>th</sup> percentile) only	<b>2.5% [2.2, 2.8] (&lt;.001)</b>	<b>2.3% [2.0, 2.6] (&lt;.001)</b>	<b>0.5% [0.1, 0.8] (0.004)</b>	<b>-0.9% [-1.1, -0.7] (&lt;.001)</b>	<b>-0.9% [-1.1, -0.7] (&lt;.001)</b>	<b>-0.4% [-0.7, -0.2] (&lt;.001)</b>
Low Medicare facilities (15 <sup>th</sup> percentile) only	<b>1.9% [1.5, 2.3] (&lt;.001)</b>	<b>1.7% [1.4, 2.1] (&lt;.001)</b>	<b>0.6% [0.5, 0.8] (&lt;.001)</b>	<b>-0.5% [-0.8, -0.4] (&lt;.001)</b>	<b>-0.6% [-0.8, -0.4] (&lt;.001)</b>	<b>-0.3% [-0.5, -0.2] (&lt;.001)</b>
High vs low Medicaid census nursing homes						
High Medicaid facilities (70 <sup>th</sup> percentile) only	<b>2.1% [1.9, 2.4] (&lt;.001)</b>	<b>2.2% [1.9, 2.4] (&lt;.001)</b>	<b>0.8% [0.7, 0.9] (&lt;.001)</b>	<b>-0.7% [-0.9, -0.5] (&lt;.001)</b>	<b>-0.8% [-0.9, -0.6] (&lt;.001)</b>	<b>-0.4% [-0.5, -0.3] (&lt;.001)</b>
Low Medicaid facilities (30 <sup>th</sup> percentile) only	<b>2.3% [2.1, 2.5] (&lt;.001)</b>	<b>2.1% [1.9, 2.3] (&lt;.001)</b>	<b>0.4% [0.3, 0.6] (&lt;.001)</b>	<b>-0.8% [-1.0, -0.6] (&lt;.001)</b>	<b>-0.7% [-0.9, -0.6] (&lt;.001)</b>	<b>-0.3% [-0.4, -0.2] (&lt;.001)</b>
High Medicaid facilities (85 <sup>th</sup> percentile) only	<b>1.9% [1.6, 2.3] (&lt;.001)</b>	<b>2.0% [1.7, 2.3] (&lt;.001)</b>	<b>0.8% [0.6, 0.9] (&lt;.001)</b>	<b>-0.8% [-1.0, -0.5] (&lt;.001)</b>	<b>-0.8% [-1.0, -0.6] (&lt;.001)</b>	<b>-0.3% [-0.4, -0.1] (&lt;.001)</b>
Low Medicaid facilities (15 <sup>th</sup> percentile) only	<b>2.2% [1.8, 2.6] (&lt;.001)</b>	<b>2.00% [1.7, 2.3] (&lt;.001)</b>	<b>0.3% [0.1, 0.6] (0.013)</b>	<b>-0.9% [-1.2, -0.6] (&lt;.001)</b>	<b>-0.9% [-1.1, -0.7] (&lt;.001)</b>	<b>-0.2% [-0.4, -0.0] (0.014)</b>
Nursing homes with high and low percentage of patients with ADRD and cognitive impairment						
High ADRD (85 <sup>th</sup> percentile)	<b>2.1% [1.9, 2.4] (&lt;.001)</b>	<b>2.3% [2.0, 2.6] (&lt;.001)</b>	<b>0.6% [0.4, 0.7] (&lt;.001)</b>	<b>-0.6% [-0.8, -0.4] (&lt;.001)</b>	<b>-0.7% [-0.9, -0.5] (&lt;.001)</b>	<b>-0.4% [-0.5, -0.2] (&lt;.001)</b>
Low ADRD (15 <sup>th</sup> percentile)	<b>2.8% [2.6, 3.1] (&lt;.001)</b>	<b>2.5% [2.2, 2.8] (&lt;.001)</b>	<b>0.9% [0.7, 1.0] (&lt;.001)</b>	<b>-1.1% [-1.3, -0.9] (&lt;.001)</b>	<b>-1.0% [-1.2, -0.9] (&lt;.001)</b>	<b>-0.6% [-0.8, -0.5] (&lt;.001)</b>

\*Each line reports findings from a separate regression model that also controls for age, gender, case mix, seasonality, and facility fixed effects. Models differ by either staffing variables (eg, salaried vs contract), or facility characteristics (eg, low quality vs high quality) and sample (full and stratified). Bolded entries are those significant at  $P < .001$ .