

Journal Pre-proof



Neither Race nor Ethnicity Impact the Mortality of Residents of Veterans Affairs
Community Living Center with COVID-19

Mayyadah H. Alabdely, MD, Sonya Kothadia, MD, Taissa Bej, MS, Brigid M. Wilson,
PhD, Sunah Song, PhD, Ukwen Akpoji, PharmD, Corinne Kowal, BS, Federico Perez,
MD, MSc, Robin L.P. Jump, MD, PhD

PII: S1525-8610(22)00831-3

DOI: <https://doi.org/10.1016/j.jamda.2022.10.020>

Reference: JMDA 4491

To appear in: *Journal of the American Medical Directors Association*

Received Date: 3 August 2022

Revised Date: 26 October 2022

Accepted Date: 30 October 2022

Please cite this article as: Alabdely MH, Kothadia S, Bej T, Wilson BM, Song S, Akpoji U, Kowal C, Perez F, Jump RLP, Neither Race nor Ethnicity Impact the Mortality of Residents of Veterans Affairs Community Living Center with COVID-19, *Journal of the American Medical Directors Association* (2022), doi: <https://doi.org/10.1016/j.jamda.2022.10.020>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Published by Elsevier Inc. on behalf of AMDA -- The Society for Post-Acute and Long-Term Care Medicine.

1 **Title:** Neither Race nor Ethnicity Impact the Mortality of Residents of Veterans Affairs
2 Community Living Center with COVID-19

3
4 **Running Title:** COVID-19 Racial Disparities absent in VA CLCs

5
6 **Authors:** Mayyadah H. Alabdely MD^a, Sonya Kothadia MD^b, Taissa Bej MS^c, Brigid M.
7 Wilson PhD^{b,c}, Sunah Song PhD^d, Ukwen Akpoji PharmD^e, Corinne Kowal BS^c, Federico Perez
8 MD, MSc^{b,c} and Robin L.P. Jump MD, PhD^{b,c*}

9
10 **Affiliations:** ^aDepartment of Medicine, University Hospitals Cleveland Medical Center,
11 Cleveland, Ohio, USA. ^bDivision of Infectious Diseases & HIV Medicine in the Department of
12 Medicine, Case Western Reserve University School of Medicine, Cleveland, Ohio. ^cGeriatric
13 Research Education and Clinical Center (GRECC), VA Northeast Ohio Healthcare System,
14 Cleveland, Ohio. ^dCleveland Institute for Computational Biology at Case Western Reserve
15 University School of Medicine, Cleveland, Ohio. ^eDepartment of Pharmacy, VA Northeast Ohio
16 Healthcare System, Cleveland, Ohio, USA.

17
18 ***Corresponding Author:** Robin L. P. Jump M.D., Ph.D., VA Pittsburgh Health Care System,
19 Geriatric Research, Education and Clinical Center (OGR-U), University Drive, Pittsburgh, PA
20 15240, USA; Telephone: 412-360-2916; Fax: 412-360-4922; e-mail: robin.jump@va.gov

21
22 ***Currently at the GRECC, VA Pittsburgh Healthcare System, Pittsburgh, Pennsylvania and the**
23 **Department of Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania.**

24
25 **Keywords:** nursing homes; healthcare disparities; SARS-CoV-2; long term care

26
27 **Funding:** This was an unfunded study supported in part by funds and facilities provided by
28 Geriatric Research Education and Clinical Centers (GRECCs) at the VA Northeast Ohio
29 Healthcare System and the VA Pittsburgh Healthcare System.

30
31 **Word count:** 288/300 words (abstract); 2060/2000 words (body); 2 tables, 1 figure (of 3 total);
32 24/30 references

33
34 **Brief Summary:** In the pre-vaccine period, increased age and comorbid conditions but not
35 frailty, race, or ethnicity were independently associated with 30-day all-cause mortality among
36 VA CLC residents with a positive COVID-19 test.

47 **Conflict of Interest Disclosures**

48 None of the authors have relevant conflicts of interest to disclose. FP and RJ have received
49 research funding from Pfizer, Merck and Accelerate. RJ has participated in advisory boards for
50 Pfizer and Merck.

51
52 **Financial support**

53 This work was supported in part by funds and facilities provided by the Geriatric Research
54 Education and Clinical Centers (GRECCs) at the VA Northeast Ohio Healthcare System and the
55 VA Pittsburgh Healthcare System. This used data from the VA COVID-19 Shared Data
56 Resource as well as resources and facilities from the Department of Veterans Affairs (VA)
57 Informatics and Computing Infrastructure (VINCI), VA HSE RES 13-457. These entities had no
58 role in the design and conduct of the study; collection, management, analysis, and interpretation
59 of the data; preparation, review, or approval of the manuscript; and decision to submit the
60 manuscript for publication. The findings and conclusions in this document are those of the
61 authors, who are responsible for its content, and do not necessarily represent the views of the VA
62 or of the United States Government.

63
64 **Author contributions**

- 65
- 66 • Study concept and design: Ukwen Akpoji, Robin Jump, Federico Perez
 - 67 • Acquisition of data: Taissa Bej, Brigid Wilson, Sunah Song
 - 68 • Analysis and interpretation of data: Taissa Bej, Corrine Kowal, Brigid Wilson
 - 69 • Drafting of the manuscript: Mayyadah Alabdely, Sonya Kothadia
 - Critical revision of the manuscript for important intellectual content: all authors

1 ABSTRACT

2 **Objectives:** COVID-19 disproportionately affected nursing home residents and people from
3 racial and ethnic minorities in the US. Nursing homes in the Veterans Affairs (VA) system,
4 termed Community Living Centers (CLCs), belong to a national managed care system. In the
5 period prior to the availability of vaccines, we examined whether residents from racial and ethnic
6 minorities experienced disparities in COVID-19 related mortality.

7 **Design:** Retrospective cohort study.

8 **Setting and Participants:** Residents at 134 VA CLCs from April 14 to December 10, 2020

9 **Methods:** We used the VA's Corporate Data Warehouse to identify VA CLC residents with a
10 positive SARS-CoV-2 PCR test during or 2 days prior to their admission and without a prior case
11 of COVID-19. We assessed age, self-reported race/ethnicity, frailty, chronic medical conditions,
12 Charlson comorbidity index, the annual quarter of the infection, and all-cause 30-day mortality.
13 We estimated odds ratios (OR) and 95% confidence intervals (CIs) of all-cause 30-day mortality
14 using a mixed-effects multivariable logistic regression model.

15 **Results:** During the study period, 1133 CLC residents had an index positive SARS-CoV-2 test.
16 Mortality at 30 days was 23% for White non-Hispanic residents, 15% for Black non-Hispanic
17 residents, 10% for Hispanic residents, and 16% for other residents. Factors associated with
18 increased 30-day mortality were age ≥ 70 , Charlson comorbidity index ≥ 6 , and a positive SARS-
19 CoV-2 test between April 14 and June 30, 2020. Frailty, Black race, and Hispanic ethnicity were
20 not independently associated with an increased risk of 30-day mortality.

21 **Conclusions and Implications:** Among a national cohort of VA CLC residents with COVID-
22 19, neither Black race nor Hispanic ethnicity had a negative impact on survival. Further research

23 is needed to determine factors within the VA healthcare system that mitigate the influence of
24 systemic racism on COVID-19 outcomes in US nursing homes.

25

Journal Pre-proof

26 INTRODUCTION

27 The COVID-19 pandemic disproportionately affected residents in nursing homes. Although
28 nursing homes provide care for <1% of the adult population in the United States, 14% of deaths
29 due to COVID-19 occurred in nursing homes.^{1,2} Prior to vaccines, nursing home residents
30 accounted for over 33% of COVID-19 related deaths.³ SARS-CoV-2 infections also
31 disproportionately affected people from racial and ethnic minorities relative to White non-
32 Hispanics. The degree to which racial and ethnic disparities, a reflection of the influences of
33 systemic racism on healthcare, were observed among nursing home residents is less clear.

34
35 Several studies that considered the period prior to the availability of an effective SARS-CoV-2
36 vaccine reported that nursing homes with higher proportions of non-White residents had higher
37 rates of COVID-19 infections and mortality.⁴⁻⁷ A systematic review of studies that included
38 facility-level characteristics concluded that nursing homes' size and community prevalence of
39 COVID-19 were among the most important factors influencing outcomes.⁸ The review also
40 noted that outcomes varied by the facility racial composition. In contrast, studies that included
41 resident-level factors found that age as well as impaired cognitive and physical function, but not
42 race or ethnicity, were associated with an increased risk of COVID-19 related mortality among
43 nursing home residents.^{9,10} Considering both facility- and resident-level characteristics, Lu *et al.*
44 assessed risk factors for COVID-19 deaths among Medicare beneficiaries aged ≥ 65 and living in
45 a nursing home. The authors reported that for-profit ownership and low health inspection ratings
46 (facility characteristics) as well as age, gender, and cognitive impairment (resident
47 characteristics) were all risk factors for COVID-19 related mortality.¹¹ Lu *et al.* proposed that

48 equal access to care, such as that provided by Medicare benefits for older persons, would
49 mitigate the impacts of systemic racism observed in prior studies of nursing home residents.

50

51 The Veterans Health Administration is a national managed care system that offers eligibility and
52 benefits based on prior military service. The Department of Veterans Affairs (VA) has 134
53 nursing homes, termed Community Living Centers (CLCs). We hypothesized that among VA
54 CLC residents with COVID-19, age, frailty, and chronic medical conditions, but not race nor
55 ethnicity, would be associated with mortality. To test our hypothesis, we conducted a
56 retrospective cohort study of VA CLC residents with documented SARS-CoV-2 infection prior
57 to the availability of COVID-19 vaccines and assessed 30-day all-cause mortality.

58

59 **METHODS**

60 **Study Design, Setting, and Data Sources.** We conducted a retrospective cohort study of
61 residents living in any of 134 VA CLCs from April 14, 2020 to December 10, 2020. On April
62 14, 2020 the VA issued a memorandum calling for widespread SARS-CoV-2 testing of all CLC
63 residents. After December 10th 2020, COVID-19 vaccines became available to VA CLC
64 residents.

65

66 We used the Veterans Affairs Informatics and Computing Infrastructure (VINCI) to access
67 clinical databases from the US Veterans Healthcare Administration (VHA). Data were extracted
68 from the VHA's Corporate Data Warehouse (CDW), the VHA's Vital Status File, and the VA
69 COVID-19 Shared Data Resource. The Institutional Review Board (IRB) at the VA [Redacted
70 for review] Healthcare System approved the study protocol.

71
72 **Case Ascertainment and Clinical Characteristics.** The cohort included all VA CLC residents
73 with an RT-PCR-based assay that was positive for SARS-CoV-2 within 2 days prior to or during
74 their admission to a VA CLC. Particularly in the early phase of the pandemic, residents may
75 have been asymptomatic or pre-symptomatic while in the CLC and subsequently found to be
76 positive while admitted to other settings within the same VA medical center (*i.e.*, acute care).
77 Therefore, residents who tested positive for SARS-CoV-2 in the week following transfer from
78 the CLC to another unit in a VA medical center were also included in the cohort. For each CLC
79 resident, only the index case (first case) was included, and individuals with a prior positive
80 SARS-CoV-2 PCR test or prior documentation of a COVID-19 case were excluded. We assessed
81 age, gender, self-reported race, self-reported ethnicity, annual quarter of the positive test, and,
82 based on *International Classification of Diseases (ICD)* and/or procedure codes, the VA Frailty
83 Index, the Charlson comorbidity index (CCI), and chronic comorbid conditions.^{12,13} All-cause
84 mortality was evaluated at 30 days following a resident's first positive SARS-CoV-2 test.

85
86 **Statistical Analysis.** Case patient characteristics were summarized by race and ethnicity.
87 Differences in mean age, frailty index, and CCI were compared using least-squares regression
88 and, when omnibus differences were detected, we performed Tukey-adjusted pair-wise
89 comparisons. Kaplan-Meier survival curves, stratified by race and ethnicity and considering
90 mortality events within 30 days of positive SARS-CoV-2 PCR test, were compared using an
91 omnibus log-rank test. A mixed-effects multivariable logistic regression model was used to
92 estimate odds ratios (OR) and 95% confidence intervals (CI) for all-cause 30-day mortality,
93 which included age, gender, race and ethnicity, VA Frailty Index, CCI, and the date of the

94 positive test (stratified by quarter from April to December 2020) as fixed effects. With the
95 mixed-effects model, we estimated random facility effects as patient outcomes within a single
96 facility may be correlated. Statistical analyses were performed using R (version 4.1.2; Vienna,
97 Austria) including functions from additional packages.¹⁴

98

99 **RESULTS**

100 After excluding patients with prior positive SARS-CoV-2 PCR tests or documented COVID-19
101 cases, we identified 1,133 CLC residents with an index positive SARS-CoV-2 PCR test between
102 April 14th and December 10th, 2020. The majority were male (1105, 98%) and their mean age
103 was 75.6 years (± 10.4) (**Table 1**). The average age of White non-Hispanic residents 77.0 years
104 (± 10.2) was greater than that of Black non-Hispanic (72.4 (± 9.7); $P < 0.001$) and Hispanic
105 residents (72.4 (± 9.7); $P < 0.001$; **Supplemental Table 1**). The mean CCI for White non-
106 Hispanic residents (4.53 (± 2.7)) was lower than that observed for Black non-Hispanic residents
107 (5.12 (± 3.1); $P = 0.021$).

108

109 All-cause 30-day mortality for CLC residents following a positive SARS-CoV-2 PCR test was
110 20% (229/1133). When stratified by race and ethnicity, all-cause mortality at 30 days was higher
111 among White non-Hispanic residents (175/758, 23%) compared to Black non-Hispanic residents
112 (36/246, 15%), Hispanic residents (4/42, 10%), and all other residents (14/87, 16%; **Figure 1**).

113 To assess the independent contribution of factors influencing mortality, we used a mixed-effects
114 multivariable logistic regression model in which CLC facility was a random effect; all other
115 variables were fixed effects. Among CLC residents with a positive SARS-CoV-2 PCR test, the
116 most notable factor impacting 30-day all-cause mortality was advanced age, increasing from an

117 OR of 5.98 (95% CI 1.407 - 25.433) for residents aged 70-79 years to an OR of 19.0 (95% CI
118 4.309 - 83.854) for residents aged ≥ 90 years relative to residents aged < 60 (**Table 2**). Residents
119 with a CCI score ≥ 6 had increased odds of 30-day mortality (OR 1.57; 95% CI 1.021 - 2.413). In
120 contrast, frailty was not associated with increased odds of mortality, and a positive test after
121 October-December 2020 was associated with decreased odds of mortality (OR 0.560; 95% CI
122 0.359 - 0.874). Of note, neither Black race nor Hispanic ethnicity (versus White-Non-Hispanic)
123 was associated with increased odds of mortality in our multivariate logistic regression model.

124

125 **DISCUSSION**

126 This retrospective study of VA CLC residents with positive SARS-CoV-2 PCR tests during the
127 pre-vaccine period found an overall all-cause 30-day mortality rate of 20%, and advanced age
128 was the resident-level factor most strongly associated with mortality. Infection early in the
129 epidemic and a high burden of comorbid medical conditions, but not frailty, were also associated
130 with increase odds of mortality. Importantly, we found that in this national cohort of VA CLC
131 residents with COVID-19, Black or Hispanic race and ethnicity were not associated with
132 increased odds of 30-day all-cause mortality.

133

134 Previous studies reported increased mortality among White compared to Black Veterans
135 hospitalized for common conditions, including pneumonia and exacerbations of congestive heart
136 failure or chronic obstructive pulmonary disease.¹⁵⁻¹⁷ In these studies, the mean age of White
137 non-Hispanic patients was greater than that of Black non-Hispanic patients. Attempts to account
138 for age and several other demographic, clinical, and social variables did not alter the findings
139 supporting similar or better outcomes for Black Veterans. Our results differ somewhat in that our

140 multivariable logistic regression model implicated advanced age as an independent risk factor for
141 mortality for CLC residents with a positive SARS-CoV-2 test. Additionally, two assessments
142 from the pre-vaccine period found that people from racial and ethnic minorities were more likely
143 to get tested for COVID-19 through the VA and were more likely to have a positive test
144 compared to White non-Hispanic people.^{18,19} Similar to our results, the rates of all-cause 30-day
145 mortality among those with positive tests did not differ by Veterans' race or ethnicity. Taken
146 together, these studies indicate that – in a healthcare system where access to care is based on
147 qualifying military service rather than employment or personal finances – people from racial and
148 ethnic minorities do not appear to have worse outcomes compared to White non-Hispanic
149 people.

150
151 The literature describing differences in outcomes prior to the availability of vaccines among
152 nursing home residents from racial and ethnic minorities is mixed. Among nursing home
153 residents with symptomatic COVID-19 infections, Panagiotou *et al.* found a lower risk of 30-day
154 all-cause mortality among non-White residents.¹⁰ Mehta *et al.* noted an increased risk of
155 hospitalization, but not death, among Black, Asian, and Hispanic or Latino nursing home
156 residents with a COVID-19 infection.⁹ Both teams indicated that risk of mortality among
157 nursing home residents with COVID-19 increased with age, impaired cognition, and limited
158 physical function. Three other studies assessed the communities in which nursing homes are
159 located and found that residents of nursing homes with the highest proportion of non-White
160 residents also had the highest rates of COVID-19 related mortality.⁴⁻⁶ Together, these findings
161 suggest that the racial segregation that continues to affect the communities surrounding nursing

162 homes has a negative influence on the health of nursing home residents from racial and ethnic
163 minorities.²⁰

164
165 The results reported here may also reflect characteristics of the VA healthcare system. While
166 most nursing homes care for people that came from the same local community, VA CLCs
167 typically have a much larger geographic catchment area and reside on a large campus that also
168 offers acute and ambulatory care services. Additionally, while VA CLCs also experienced
169 staffing shortages and challenges related to obtaining adequate testing supplies and personal
170 protective equipment, being part of an integrated healthcare system with a unionized nursing
171 staff may have mitigated some of these shortfalls.²¹ Specifically, within each VA institution, staff
172 from other areas could be reassigned to care for CLC residents. Additionally, regional networks
173 within the VA system, termed Veterans Integrated Service Networks (VISNs), could help direct
174 personnel protective equipment (PPE) to the VA medical centers with the greatest need.

175
176 Our study has important limitations. First, our cohort consists of VA healthcare users, a
177 predominantly male and White non-Hispanic population with a high burden of comorbidities.^{22,23}

178 While all-cause mortality at 30 days among VA CLC residents with a positive SARS-CoV-2
179 PCR test was similar to that reported for residents of community nursing homes,^{9,10} other
180 differences between Veteran and non-Veteran population, such as the higher proportion of males,
181 affects the comparability of our findings to those of studies among nursing home residents in
182 community settings.²⁴ Second, our cohort of VA CLC residents was relatively small; especially
183 cases among veterans who identified as Hispanics or Other were very few, and may reflect
184 perceived ambiguity in these classifications. While extending the study period would have

185 permitted a larger cohort, we restricted our analysis to the period during which CLC residents
186 were routinely tested for SARS-CoV-2 and prior to the available of effective vaccines. These
187 restrictions limited the statistical power to detect differences between groups, especially for
188 people included in the Hispanic and Other categories.

189

190 **CONCLUSIONS AND IMPLICATIONS**

191 Among residents of VA CLCs with a positive SARS-CoV-2 PCR test, Black and Hispanic
192 residents did not experience higher all-cause 30-day mortality compared to White non-Hispanic
193 residents. Rather, factors such as advanced age, high comorbidity burden, conditions and
194 infection early in the epidemic were independently associated with increased mortality. The VA
195 is the largest integrated healthcare system in the US and differs from other healthcare
196 organizations in that eligibility is based on prior military service, rather than age, socioeconomic
197 status, or location. Greater understanding of aspects of the VA healthcare system that have the
198 potential to abate racial and ethnic disparities among VA CLC residents with COVID-19 may
199 help inform efforts to mitigate systemic racism on outcomes across non-VA nursing homes and
200 other healthcare sectors.

201 **REFERENCES**


- 202 1. National Center for Health Statistics (U.S.), ed. *Long-Term Care Providers and Services*
203 *Users in the United States, 2015-2016*. U.S. Department of Health and Human Services,
204 Centers for Disease Control and Prevention, National Center for Health Statistics; 2019.
- 205 2. NVSS - Provisional Death Counts for COVID-19 - Executive Summary.
- 206 3. Times TNY. More Than One-Third of U.S. Coronavirus Deaths Are Linked to Nursing Homes.
207 *The New York Times*. [https://www.nytimes.com/interactive/2020/us/coronavirus-nursing-](https://www.nytimes.com/interactive/2020/us/coronavirus-nursing-homes.html)
208 [homes.html](https://www.nytimes.com/interactive/2020/us/coronavirus-nursing-homes.html). Published June 27, 2020. Accessed March 10, 2021.
- 209 4. Gorges RJ, Konetzka RT. Factors Associated With Racial Differences in Deaths Among
210 Nursing Home Residents With COVID-19 Infection in the US. *JAMA Netw Open*.
211 2021;4(2):e2037431.
- 212 5. Cai S, Yan D, Intrator O. COVID-19 Cases and Death in Nursing Homes: The Role of Racial
213 and Ethnic Composition of Facilities and Their Communities. *J Am Med Dir Assoc*.
214 2021;22(7):1345-1351.
- 215 6. Li Y, Cen X, Cai X, Temkin-Greener H. Racial and Ethnic Disparities in COVID-19 Infections
216 and Deaths Across U.S. Nursing Homes. *J Am Geriatr Soc*. n/a(n/a).
- 217 7. Kumar A, Roy I, Karmarkar AM, et al. Shifting US Patterns of COVID-19 Mortality by Race
218 and Ethnicity From June–December 2020. *J Am Med Dir Assoc*. 2021;22(5):966-970.e3.
- 219 8. Konetzka RT, White EM, Pralea A, Grabowski DC, Mor V. A systematic review of long-term
220 care facility characteristics associated with COVID-19 outcomes. *J Am Geriatr Soc*.
221 2021;69(10):2766-2777.
- 222 9. Mehta HB, Li S, Goodwin JS. Risk Factors Associated With SARS-CoV-2 Infections,
223 Hospitalization, and Mortality Among US Nursing Home Residents. *JAMA Netw Open*.
224 2021;4(3):e216315.
- 225 10. Panagiotou OA, Kosar CM, White EM, et al. Risk Factors Associated With All-Cause 30-Day
226 Mortality in Nursing Home Residents With COVID-19. *JAMA Intern Med*. 2021;181(4):439-
227 448.
- 228 11. Lu Y, Jiao Y, Graham DJ, et al. Risk Factors for COVID-19 Deaths Among Elderly Nursing
229 Home Medicare Beneficiaries in the Pre-vaccine Period. *J Infect Dis*. 2022;225(4):567-577.
- 230 12. Cheng D, DuMontier C, Yildirim C, et al. Updating and Validating the U.S. Veterans Affairs
231 Frailty Index: Transitioning From ICD-9 to ICD-10. *J Gerontol A Biol Sci Med Sci*.
232 2021;76(7):1318-1325.

- 233 13. Quan H, Sundararajan V, Halfon P, et al. Coding algorithms for defining comorbidities in
234 ICD-9-CM and ICD-10 administrative data. *Med Care*. 2005;43(11):1130-1139.
- 235 14. R Core Team. R: A language and environment for statistical computing. Published online
236 20210101.
- 237 15. Deswal A, Petersen NJ, Soucek J, Ashton CM, Wray NP. Impact of race on health care
238 utilization and outcomes in veterans with congestive heart failure. *J Am Coll Cardiol*.
239 2004;43(5):778-784.
- 240 16. Vaughan Sarrazin M, Cannon KT, Rosenthal GE, Kaldjian LC. Racial Differences in Mortality
241 Among Veterans Hospitalized for Exacerbation of Chronic Obstructive Pulmonary Disease.
242 *J Natl Med Assoc*. 2009;101(7):656-662.
- 243 17. Silva GC, Jiang L, Gutman R, et al. Racial/Ethnic Differences in 30-Day Mortality for Heart
244 Failure and Pneumonia in the Veterans Health Administration Using Claims-based, Clinical,
245 and Social Risk-adjustment Variables. *Med Care*. 2021;59(12):1082-1089.
- 246 18. Ferguson JM, Abdel Magid HS, Purnell AL, Kiang MV, Osborne TF. Differences in COVID-19
247 Testing and Test Positivity Among Veterans, United States, 2020. *Public Health Rep*.
248 2021;136(4):483-492.
- 249 19. Rentsch CT, Kidwai-Khan F, Tate JP, et al. Patterns of COVID-19 testing and mortality by
250 race and ethnicity among United States veterans: A nationwide cohort study. *PLOS Med*.
251 2020;17(9):e1003379.
- 252 20. Shippee TP, Fabius CD, Fashaw-Walters S, et al. Evidence for Action: Addressing Systemic
253 Racism Across Long-Term Services and Supports. *J Am Med Dir Assoc*. 2022;23(2):214-219.
- 254 21. Review of Veterans Health Administration's COVID-19 Response and Continued Pandemic
255 Readiness. :46.
- 256 22. Agha Z, Lofgren RP, VanRuiswyk JV, Layde PM. Are patients at Veterans Affairs medical
257 centers sicker? A comparative analysis of health status and medical resource use. *Arch*
258 *Intern Med*. 2000;160(21):3252-3257.
- 259 23. Dursa EK, Barth SK, Bossarte RM, Schneiderman AI. Demographic, Military, and Health
260 Characteristics of VA Health Care Users and Nonusers Who Served in or During Operation
261 Enduring Freedom or Operation Iraqi Freedom, 2009-2011. *Public Health Rep*.
262 2016;131(6):839-843.
- 263 24. Harris-Kojetin L, Lendon J, Rome V, Valverde R, Caffrey C. Long-Term Care Providers and
264 services users in the United States, 2015-2016. *Vital Health Stat Ser 3 Anal Epidemiol Stud*
265 *US Dept Health Hum Serv Public Health Serv Natl Cent Health Stat*. 2019;(43).

266

267 **FIGURE LEGEND**

268 **Figure 1:** Survival Curves of VA Community Living Center Residents with a Positive SARS-
 269 CoV-2 PCR test. White = grey continuous line; Black = black continuous curve; Hispanic =
 270 black dotted curve; Other = grey dotted curve. Other includes American Indian or Alaska
 271 Native, Asian, Multiple, Unknown, Missing, or Declined. Note that the y-axis starts at 0.5.
 272 Comparing the survival curves with an omnibus log-rank test yielded a P-value of 0.01.

Number at risk

	0	10	20	30
White, non-Hispanic	758	699	620	583
Black, non-Hispanic	246	239	221	210
Hispanic	42	39	38	38
Other	87	81	76	73

273

274 TABLES

275 **Table 1:** VA CLC Residents with a Positive SARS-CoV-2 PCR Test, April 14th – December
276 10th, 2020

Characteristics	All Cases N = 1133	White Non- Hispanic n = 758	Black Non- Hispanic n = 246	Hispanic n = 42	Other ^a n = 87
Age, mean (\pm SD)	75.6 (\pm 10.4)	77.0 (\pm 10.2)	72.4 (\pm 9.7)	70.6 (\pm 11.9)	75.4 (\pm 11.3)
Male, No. (%)	1084 (97%)	719 (98%)	241 (96%)	41 (98%)	83 (99%)
Frailty Index (\pm SD)	0.41 (\pm 0.1)	0.42 (\pm 0.1)	0.40 (\pm 0.1)	0.40 (\pm 0.1)	0.39 (\pm 0.1)
Charlson Comorbidity Index, mean (\pm SD)	4.65 (\pm 2.8)	4.53 (\pm 2.7)	5.12 (\pm 3.1)	4.60 (\pm 2.9)	4.43 (\pm 2.3)
Chronic Medical Conditions					
Dementia	681 (60%)	459 (61%)	145 (59%)	25 (60%)	52 (60%)
Diabetes	578 (51%)	368 (49%)	142 (58%)	23 (55%)	45 (52%)
Chronic pulmonary disease	447 (39%)	314 (41%)	85 (35%)	14 (33%)	34 (39%)
Stroke	394 (35%)	234 (31%)	110 (45%)	18 (43%)	32 (37%)
Peripheral vascular disease	374 (33%)	250 (33%)	75 (30%)	18 (43%)	31 (36%)
Renal disease	342 (30%)	229 (30%)	75 (30%)	9 (21%)	29 (33%)
Congestive heart failure	339 (30%)	235 (31%)	73 (30%)	9 (21%)	22 (25%)
Cancer	262 (23%)	175 (23%)	60 (24%)	10 (24%)	17 (20%)
Liver disease	153 (14%)	80 (11%)	53 (22%)	10 (24%)	10 (11%)
Paralysis	102 (9%)	61 (8%)	29 (12%)	4 (10%)	8 (9%)
Myocardial infarction	100 (9%)	79 (10%)	14 (6%)	1 (2%)	6 (7%)
Peptic ulcer disease	58 (5%)	38 (5%)	15 (6%)	2 (5%)	3 (3%)
Rheumatic disease	28 (2%)	17 (2%)	5 (2%)	3 (7%)	3 (3%)
HIV	8 (1%)	0 (0%)	7 (3%)	1 (2%)	0 (0%)

277 ^a Other includes American Indian or Alaska Native, Asian, Multiple, Unknown, Missing, or
278 Declined

279
280
281
282
283
284


285 **Table 2.** Mixed-effects Multivariable Logistic Regression Odds Ratio for 30-Day All-Cause
 286 Mortality Among VA CLC Residents with a Positive SARS-CoV-2 PCR Test Between April
 287 14th and December 10th, 2020, CLC facility as random effect, fixed effects shown

Characteristics	Adjusted Odds Ratio	95% Confidence Interval	P-value ^a
Age (reference: <60 years)			
60 – 69 years	2.790	(0.629,12.382)	0.177
70 – 79 years	5.981	(1.407,25.433)	0.015
80 – 89 years	9.559	(2.215,41.254)	0.002
≥ 90 years	19.008	(4.309,83.854)	<0.0001
Male gender (reference: female)	0.546	(0.195,1.535)	0.251
Race & Ethnicity (reference: White, non-Hispanic)			
Black, non-Hispanic	0.720	(0.466,1.111)	0.138
Hispanic	0.473	(0.156,1.439)	0.187
Other ^b	0.726	(0.384,1.371)	0.323
Frailty (reference: non- or pre-frail)			
Mild frailty	1.267	(0.567,2.832)	0.565
Moderate frailty	1.174	(0.539,2.558)	0.687
Severe frailty	1.266	(0.581,2.758)	0.552
Charlson comorbidity index (reference: <4)			
4-5	1.371	(0.900,2.089)	0.142
≥6	1.570	(1.021,2.413)	0.040
Time (reference: April 14th – June 30th 2020)			
July 1st – September 30th 2020	0.687	(0.412,1.144)	0.149
October 1st – December 14th 2020	0.560	(0.359,0.874)	0.011

288 ^aP-values < 0.05 are considered statistically significant; these values are noted in **bold text**
 289 within the table.

290 ^bOther includes American Indian or Alaska Native, Asian, Multiple, Unknown, Missing, or
 291 Declined

292



Number at risk

	0	10	20	30
White, non-Hispanic	758	699	620	583
Black, non-Hispanic	246	239	221	210
Hispanic	42	39	38	38
Other	87	81	76	73